



This is a repository copy of *Survival outcomes in gastrostomy fed stroke patients*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/134938/>

Version: Accepted Version

Proceedings Paper:

Mallouppa, E., Naqvi, A., Kurien, M. orcid.org/0000-0002-4227-9500 et al. (2 more authors) (2016) Survival outcomes in gastrostomy fed stroke patients. In: International Journal of Stroke. UK Stroke Forum 2016, 28-30 Nov 2016, Liverpool, UK. SAGE Publications , S36-S36.

<https://doi.org/10.1177/1747493016669275>

© 2016 World Stroke Organization. This is an author produced version of a paper subsequently published in International Journal of Stroke. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Survival outcomes in gastrostomy fed stroke patients

E Mallouppa¹, A Naqvi², M Kurien³, G Pratt² and D Sanders³

¹Geriatric Medicine, Chesterfield Royal Hospital NHS Foundation Trust, Chesterfield, UK

²Stroke Department, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

³Academic Unit of Gastroenterology, Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, UK

Introduction

Dysphagia following acute stroke is common. It is associated with increased morbidity and mortality and longer hospital length of stay. There is little evidence on survival outcomes in these patients.

Method

Retrospective observational study looking at 153 consecutive patients with acute stroke who underwent gastrostomy insertion. Survival was assessed using Kaplan Meier survival curves. Associations between survival and age, sex and timing of gastrostomy insertion were analysed using log-rank. Multivariate analysis was performed with Cox regression proportional hazards.

Results

At 2 years 21/153 (13.7%) patients were alive. Median survival in patients 80 or less was 270 days compared to 98 days in the over 80s (log-rank p-value <0.0001). Median survival for women was 158 days compared to 97 days in men (log-rank p-value: 0.014). There was no association between timing of gastrostomy insertion and survival (median survival for insertion within 28 days 158 days vs 139 days for insertion after 28 days p-value 0.997). On multivariable analysis, increasing age (as a continuous variable) and male gender were independently associated with a worse prognosis (HR 1.07 and 1.63 respectively).

Conclusion

2 year survival in these patients is poor. Advancing age and male sex are associated with a worse prognosis whereas timing of gastrostomy insertion is not related to outcome. This data supports findings from previous studies, which fail to show an advantage of early gastrostomy feeding. This study is important as it allows clinicians to provide patients and relatives with realistic information for future outcomes to allow informed decisions with regards to gastrostomy feeding.